

BIOGUARD

HIGH-EFFICIENCY FOAMING SANITIZER

BIOGUARD is a Peracetic acid and Hydrogen Peroxide based sanitizer used for sanitizing agricultural and horticultural premises such as hog barns and poultry housing facilities, hatcheries and livestock trucks and trailers on all surfaces and equipment.

BIOGUARD'S wide spectrum acts on most bacteria, mould and yeast.

DO NOT STORE WITH FOOD PRODUCTS.

FEATURES AND BENEFITS

- · High foaming liquid
- · Strong oxidizer
- Biodegradable
- · Low reactivity with organic matter
- Effective in cold and hard water
- No cell resistance build-up
- High lipid solubility
- Freedom from deactivation by catalase and peroxidase
- Extremely efficient and fast acting at low concentration



DIRECTION FOR USE

Equipment to be treated

Agricultural and horticultural equipment and premises: trucks, ceilings, walls, floors.

Use procedure

- 1. Remove all animals and feed.
- 2. Remove all dirt, manure, debris from floor or surface.
- 3. Pre-rinse with water to pre-clean. (High pressure if possible).
- 4. Clean with an alkaline cleaner.
- 5. Rinse with water. (High pressure if possible).
- 6. Apply **BIOGUARD** to sanitize the cleaned surface by using a foaming applicator, or by fogging.
- 7. Allow the surface to air for 4 hours before admitting livestock.

Concentration

600 ppm to 1000 ppm (1.0 % to 1.6 %).

Contact time

Minimum 10 minutes. Do not house livestock or use equipment until treatment has been absorbed and set to dry.

Temperature

Between 4°C and 40°C (39 and 105°F).

Rinsing

All food contact surfaces, treated feed racks, mangers, troughs, automatic feeders, fountains and waterers must be thoroughly rinsed with potable prior to reuse.



BIOGUARD

HIGH-EFFICIENCY FOAMING SANITIZER



BACTERIAL INACTIVATION BY COMMON BIOCIDAL

Bacterium	Glutaraldehyde 2%	Formaldehyde 8%	Phenol 5%	Cu/Asc 0.1%	Hypochlorite 0.05%	Peroxide 10%	Peracetic 0.03%
B.cereus	> 5.0(2)	> 5.0(2)	> 5.0(2)	> 5.0(2)	> 5.0(2)	> 5.0(2)	> 5.0(2)
C.perfringens	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.3(2)	0.14±0.05(2)	> 6.3(2)	4.1±0.1(2)
E.coli	> 6.9(2)	> 6.9(2)	> 6.9(2)	6.3±0.8(2)	$6.2 \pm 0.9(2)$	> 6.9(2)	> 6.9(2)
L.monocytogenes	> 6.1(2)	> 6.1(1)	> 6.1(2)	> 6.1(1)	> 6.1(2)	> 6.1(2)	> 6.1(1)
P. aeruginosa	3.8±0.2(2)	> 6.1(3)	5.8±0.6(3)	5.6±0.9(3)	1.3±0.1(2)	> 6.1(3)	5.0±1.6(3)
S.typhimurium	> 6.4(3)	> 6.2(3)	> 6.4(3)	> 6.4(3)	4.1±1.3(2)	> 6.4(3)	> 6.4(3)
S.sonnei	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.1(1)	> 6.3(2)	> 6.3(2)	> 6.3(2)
S.aureus	> 6.5(3)	> 6.3(3)	> 6.3(3)	5.5±1.2(3)	4.8±1.8(2)	5.6±0.7(3)	6.6±03(3)
S. epidermidis	> 6.3(2)	5.9±1.1(3)	> 6.3(2)	5.1±0.1(2)	6.3±0.4(3)	> 6.3(3)	> 6.3(3)
V.cholerae	> 6.4(2)	> 6.4(2)	> 6.4(2)	> 6.4(2)	> 6.4(2)	> 6.4(2)	> 6.4(2)
V.parahaemolyticus	> 6.2(1)	> 6.2(2)	> 6.2(2)	> 6.2(2)	> 6.2(2)	> 6.2(2)	> 6.2(2)
V.vulniticus	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.3(2)	> 6.3(2)
Y.enterocolitica	> 6.8(2)	> 6.8(2)	> 6.8(2)	> 6.8(2)	> 6.8(2)	> 6.8(2)	> 6.8(2)

Calculated as $-\log$ (Td / Tw) where Td is the titer of bacteria surviving 30 min exposure at 20°C to a given disinfectant, and Tw is the titer of bacteria exposed under the same condition to water results are expressed either as the limit of detection when no surviving colonies were obtained or as $x \pm s$ (n) where n is the number of replicate experiments.

From Sagripanti J-L,Eklund CA, TrstPA et al.Comparative sensitivity of 13 species of pathogenic bacteria to seven germicide. Am J Infect Control 1997;25:335-339

Disinfection, Sterilization, and Preservation EDITOR SEYMOUR S. BLOCK FIFTH EDITION

PHYSICAL PROPERTIES

Appearence

Odour

Specific gravity @ 20°C

Viscosity

pН

Flash point (TCC)

Water solubility

Auto ignition temperature

Chlorine

Boiling point

Freezing point

Foaming tendency

Concentration monitoring techniques

Rinsing

Clear, colorless liquid					
Pungent vinegar					
1.10					
Not available					
1.2					
None					
Complete					
None					
None					
100°C					

INCOMPATIBILITY

Chlorinated products, alkalis, metals, organic materials

WAREHOUSING PRECAUTIONS

Keep containers tightly closed in a dry, cool and well-ventilated place

AVAILABLE SIZES							
	20L	205L					







IMPORTANT

Before using **BIOGUARD**, always be sure to read and follow precautions and directions for use appearing on the product's container label, and on the safety data sheet (S.D.S).

Electrical conductivity, titration

FIRST AID MEASURES

Eye contact

Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician.

-5°C

High

Excellent

Skin contact

Remove/Take off immediately all contaminated clothing. Rinse skin with water/ shower. Wash contaminated clothing before reuse.

Inhalation

Remove person to fresh air and keep at rest in a position comfortable for breathing. Immediately call a POISON CENTER or doctor/physician.

Ingestion

Do not induce vomiting. Drink 1 or 2 glasses of water. Call a physician or Poison Control Centre immediately. Never give anything by mouth to an unconscious person.

PERSONAL PROTECTION









MONTREAL

10,900 Secant Street Anjou, QC H1J 1S5 Tel.: 514-355-4660 Toll free : 1 800-361-8927

EDMONTON

11411-160 Street Edmonton, Alberta T5M 3T7 Tel.: 780-454-3919 Toll free : 1 866-454-3919